

NDA 2 2024

LIVE

MATHS

REVISION

CLASS 9



NAVJYOTI SIR



16 August 2024 Live Classes Schedule

8:00AM	16 AUGUST 2024 DAILY CURRENT AFFAIRS	RUBY MA'AM
9:00AM	16 AUGUST 2024 DAILY DEFENCE UPDATES	DIVYANSHU SIR

SSB INTERVIEW LIVE CLASSES

9:00AM	COMPLETE PSYCH TEST	ANURADHA MA'AM
--------	---------------------	----------------

NDA 2 2024 LIVE CLASSES

11:00AM	GK - GEOGRAPHY REVISION - CLASS 3	RUBY MA'AM
✓ 1:00PM	MATHS REVISION - CLASS 3	NAVJYOTI SIR
2:00PM	CHEMISTRY REVISION - CLASS 2	SHIVANGI MA'AM
5:30PM	ENGLISH - REVISION - CLASS 5	ANURADHA MA'AM

CDS 2 2024 LIVE CLASSES

11:00AM	GK - GEOGRAPHY REVISION - CLASS 3	RUBY MA'AM
2:00PM	CHEMISTRY REVISION - CLASS 2	SHIVANGI MA'AM
3:00PM	MATHS REVISION - CLASS 9	NAVJYOTI SIR
5:30PM	ENGLISH - REVISION - CLASS 5	ANURADHA MA'AM



REVISION TOPICS :

- **Matrices and Determinants**

Q) Consider the following statements:

✓ 1. If $\det A = 0$, then $\det(\operatorname{adj} A) = 0$

✓ 2. If A is non-singular, then $\det(A^{-1}) = (\det A)^{-1}$

(a) 1 only

(b) 2 only

✓ (c) Both 1 and 2

(d) Neither 1 nor 2

$$\textcircled{1} \det(\operatorname{adj} A) = |A|^{n-1} = (0)^{n-1} = \underline{0}$$

$$\textcircled{2} |A| \neq 0,$$

$$\checkmark \det(A^{-1}) = \frac{1}{(\det A)}$$

Q) Consider the following statements:

1. If $\det A = 0$, then $\det(\operatorname{adj} A) = 0$

2. If A is non-singular, then $\det(A^{-1}) = (\det A)^{-1}$

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Ans: (c)

Q) If $l + m + n = 0$, then the system of equations

$$-2x + y + z = l$$

$$x - 2y + z = m$$

$$x + y - 2z = n$$

$$A = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix} \quad B = \begin{bmatrix} l \\ m \\ n \end{bmatrix}$$

has

(a) a trivial solution

(b) no solution

(c) a unique solution

(d) infinitely many solutions

$\det(A) = 0$ &

① $|A| = -2(3) - 1(-3) + 1(3) = -6 + 3 + 3 = 0$

$(\text{adj}^{\circ}A)B = 0$,

② $(\text{adj}^{\circ}A)(B) = \begin{bmatrix} 3 & 3 & 3 \\ 3 & 3 & 3 \\ 3 & 3 & 3 \end{bmatrix} \begin{bmatrix} l \\ m \\ n \end{bmatrix} = \begin{bmatrix} 3(l+m+n) \\ 3(l+m+n) \\ 3(l+m+n) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

Q) If $l + m + n = 0$, then the system of equations

$$-2x + y + z = l$$

$$x - 2y + z = m$$

$$x + y - 2z = n$$

has

- (a) a trivial solution (b) no solution
(c) a unique solution (d) infinitely many solutions

Ans: (d)

Q) Consider the following statements in respect of symmetric matrices A and B

1. AB is symmetric. ✗
2. $A^2 + B^2$ is symmetric. ✓

A

$A' = A$ (A' is transpose of A)

Which of the above statement(s) is/are correct?

- (a) 1 only (b) 2 only ✓
(c) Both 1 and 2 (d) Neither 1 nor 2

① $(AB)' = AB$ (if AB has to be symmetric)

LHS = $(AB)' = B'A' \neq$ RHS

② $A^2 + B^2$ = $(A^2 + B^2)'$ (if $A^2 + B^2$ has to be symmetric)

RHS = $(A^2)' + (B^2)' = A^2 + B^2 =$ LHS ✓

Q) Consider the following statements in respect of symmetric matrices A and B

1. AB is symmetric.
2. $A^2 + B^2$ is symmetric.

Which of the above statement(s) is/are correct?

- | | |
|------------------|---------------------|
| (a) 1 only | (b) 2 only |
| (c) Both 1 and 2 | (d) Neither 1 nor 2 |

Ans: (b)

Q) If $A = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix}$, where ω is cube root of unity, then what is

A^{100} equal to?

(a) A

(b) $-A$

(c) Null matrix

(d) Identity matrix

$$A^2 = A \cdot A = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix} \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix} = \begin{bmatrix} \omega^2 & 0 \\ 0 & \omega^2 \end{bmatrix}$$

$$A^3 = A^2 \cdot A = \begin{bmatrix} \omega^2 & 0 \\ 0 & \omega^2 \end{bmatrix} \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix} = \begin{bmatrix} \omega^3 & 0 \\ 0 & \omega^3 \end{bmatrix}$$

$$A^n = \begin{bmatrix} \omega^n & 0 \\ 0 & \omega^n \end{bmatrix}$$

(OR)

$$A = \omega \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$A^2 = \left(\omega \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \left(\omega \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right)$$

$$= \omega^2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \omega^2 & 0 \\ 0 & \omega^2 \end{bmatrix}$$

$$A^{100} = \omega^{100} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \omega^{100} & 0 \\ 0 & \omega^{100} \end{bmatrix} = \begin{bmatrix} \omega^1 & 0 \\ 0 & \omega^1 \end{bmatrix} = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix} = \underline{\underline{A}}$$

Q) If $A = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix}$, where ω is cube root of unity, then what is

A^{100} equal to?

- (a) A (b) $-A$
(c) Null matrix (d) Identity matrix

Ans: (a)

Q) A matrix X has $(a + b)$ rows and $(a + 2)$ columns; and a matrix Y has $(b + 1)$ rows and $(a + 3)$ columns. If both XY and YX exist, then what are the values of a, b respectively?

(a) 3, 2

(b) 2, 3

(c) 2, 4

(d) 4, 3

XY existing

$$a + 2 = b + 1$$

$$a - b = -1$$

$$\underline{\underline{a = 2}}$$

YX existing

$$a + 3 = a + b$$

$$\underline{\underline{b = 3}}$$

Q) A matrix X has $(a + b)$ rows and $(a + 2)$ columns; and a matrix Y has $(b + 1)$ rows and $(a + 3)$ columns. If both XY and YX exist, then what are the values of a, b respectively?

(a) 3, 2

(b) 2, 3

(c) 2, 4

(d) 4, 3

Ans: (b)

Q) If $f(x) = \begin{vmatrix} 1 + \sin^2 x & \cos^2 x & 4 \sin 2x \\ \sin^2 x & 1 + \cos^2 x & 4 \sin 2x \\ \sin^2 x & \cos^2 x & 1 + 4 \sin 2x \end{vmatrix}$

What is the maximum value of $f(x)$?

- (a) 2
- (b) 4
- (c) 6
- (d) 8

$C_1 \rightarrow C_1 + C_2 + C_3$

$$\begin{vmatrix} 1 + \sin^2 x + \cos^2 x + 4 \sin 2x & \cos^2 x & 4 \sin 2x \\ 1 + \sin^2 x & 1 + \cos^2 x & 4 \sin 2x \\ \sin^2 x & \cos^2 x & 1 + 4 \sin 2x \end{vmatrix}$$

NDA 2 2024 - REVISION - MATHS - CLASS 9

$$(2 + 4\sin 2x) \left| \begin{array}{cc|c} 1 & \underline{\cos^2 x} & \underline{4\sin 2x} \\ 1 & \underline{1 + \cos^2 x} & \underline{4\sin 2x} \\ 1 & \cos^2 x & 1 + 4\sin 2x \end{array} \right|$$

$$R_2 \rightarrow R_2 - R_1, \quad R_3 \rightarrow R_3 - R_2$$

$$(2 + 4\sin 2x) \left| \begin{array}{cc|c} 1 & \cos^2 x & 4\sin 2x \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{array} \right|$$

$$f(x) = 2(1 + 2\sin 2x)(1)(1 - 0) = \underline{2 + 4\sin 2x}$$

max. when $\sin 2x = 1$

$$2 + 4(1) = \textcircled{6}$$

Q) If $f(x) = \begin{vmatrix} 1 + \sin^2 x & \cos^2 x & 4 \sin 2x \\ \sin^2 x & 1 + \cos^2 x & 4 \sin 2x \\ \sin^2 x & \cos^2 x & 1 + 4 \sin 2x \end{vmatrix}$

What is the maximum value of $f(x)$?

- (a) 2 (b) 4
(c) 6 (d) 8

Ans: (c)

Q) For a square matrix A , which of the following properties hold?

1. $(A^{-1})^{-1} = A$

2. $\det(A^{-1}) = \frac{1}{\det A}$

3. $(\lambda A)^{-1} = \lambda A^{-1}$, where λ is a scalar

Select the correct answer using the code given below.

- (a) 1 and 2 (b) 2 and 3 (c) 1 and 3 (d) 1, 2 and 3

Q) For a square matrix A , which of the following properties hold?

1. $(A^{-1})^{-1} = A$

2. $\det(A^{-1}) = \frac{1}{\det A}$

3. $(\lambda A)^{-1} = \lambda A^{-1}$, where λ is a scalar

Select the correct answer using the code given below.

- (a) 1 and 2 (b) 2 and 3 (c) 1 and 3 (d) 1, 2 and 3

Ans: (d)

Q) The system of equations

$$2x + y - 3z = 5$$

$$3x - 2y + 2z = 5 \text{ and } 5x - 3y - z = 16$$

(a) is inconsistent

(b) is consistent, with a unique solution ✓

(c) is consistent, with infinitely many solutions

(d) has its solution lying along X-axis in three-dimensional space

$$A = \begin{bmatrix} 2 & 1 & -3 \\ 3 & -2 & 2 \\ 5 & -3 & -1 \end{bmatrix} \Rightarrow |A| = 2(8) - 1(-3-10) - 3(-9+10)$$

$$= \underline{16 + 13 - 3} = 16 + 10 = 26$$

As $|A| \neq 0$, so consistent with unique solution.

Q) The system of equations

$$2x + y - 3z = 5$$

$$3x - 2y + 2z = 5 \text{ and } 5x - 3y - z = 16$$

- (a) is inconsistent
- (b) is consistent, with a unique solution
- (c) is consistent, with infinitely many solutions
- (d) has its solution lying along X-axis in three-dimensional space

Ans: (b)

Q) If $\Delta = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$

then what is

$\begin{vmatrix} 3d + 5g & 4a + 7g & 6g \\ 3e + 5h & 4b + 7h & 6h \\ 3f + 5i & 4c + 7i & 6i \end{vmatrix}$ equal to?

- (a) Δ ✓
- (b) 7Δ ✓
- (c) 72Δ ✓
- (d) -72Δ ✓

$6 \begin{vmatrix} \underline{3d+5g} & \underline{4a+7g} & \underline{6g} \\ \underline{3e+5h} & \underline{4b+7h} & \underline{6h} \\ \underline{3f+5i} & \underline{4c+7i} & \underline{6i} \end{vmatrix}$

$\Rightarrow C_1 \rightarrow C_1 - 5C_3 \quad C_2 \rightarrow C_2 - 7C_3$

$6 \begin{vmatrix} 3d & 4a & g \\ 3e & 4b & h \\ 3f & 4c & i \end{vmatrix} = 6 \times 3 \times 4 \begin{vmatrix} d & a & g \\ e & b & h \\ f & c & i \end{vmatrix}$

$= -72 \begin{vmatrix} a & d & g \\ b & e & h \\ c & f & i \end{vmatrix} = -72 \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$

$= -72\Delta$

Q) If $\Delta = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$

then what is

$$\begin{vmatrix} 3d + 5g & 4a + 7g & 6g \\ 3e + 5h & 4b + 7h & 6h \\ 3f + 5i & 4c + 7i & 6i \end{vmatrix} \text{ equal to?}$$

(a) Δ

(b) 7Δ

(c) 72Δ

(d) -72Δ

Ans: (d)

Q) If $A = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$,

where $a \in \mathbb{N}$, then what is $A^{100} - A^{50} - 2A^{25}$ equal to?

- (a) $-2I$ (b) $-I$
 (c) $2I$ (d) I

where I is the identity matrix.

$$A^{100} = \begin{bmatrix} 1 & 100a \\ 0 & 1 \end{bmatrix}$$

$$A^{50} = \begin{bmatrix} 1 & 50a \\ 0 & 1 \end{bmatrix}$$

$$2A^{25} = \begin{bmatrix} 2 & 50a \\ 0 & 2 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2a \\ 0 & 1 \end{bmatrix}$$

$$A^3 = A^2 \cdot A = \begin{bmatrix} 1 & 2a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 3a \\ 0 & 1 \end{bmatrix}$$

$$A^{100} - A^{50} - 2A^{25} = \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} = -2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = -2I$$

(2)

Q) If $A = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$,

where $a \in \mathbb{N}$, then what is
 $A^{100} - A^{50} - 2A^{25}$ equal to?

- (a) $-2I$ (b) $-I$
(c) $2I$ (d) I

where I is the identity matrix.

Ans: (a)

Q) If A and B are square matrices of order 2 such that $\det(AB) = \det(BA)$, then which one of the following is correct?

- (a) A must be a unit matrix
- (b) B must be a unit matrix
- (c) Both A and B must be unit matrices
- (d) A and B need not be unit matrices

Q) If A and B are square matrices of order 2 such that $\det(AB) = \det(BA)$, then which one of the following is correct?

- (a) A must be a unit matrix
- (b) B must be a unit matrix
- (c) Both A and B must be unit matrices
- (d) A and B need not be unit matrices

Ans: (d)

Q) If α and β are the roots of the equation $1 + x + x^2 = 0$,

then the matrix product $\begin{bmatrix} 1 & \beta \\ \alpha & \alpha \end{bmatrix} \begin{bmatrix} \alpha & \beta \\ 1 & \beta \end{bmatrix}$ is equal to

(a) $\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$

(b) $\begin{bmatrix} -1 & -1 \\ -1 & 2 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$

(d) $\begin{bmatrix} -1 & -1 \\ -1 & -2 \end{bmatrix}$

Q) If α and β are the roots of the equation $1 + x + x^2 = 0$,

then the matrix product $\begin{bmatrix} 1 & \beta \\ \alpha & \alpha \end{bmatrix} \begin{bmatrix} \alpha & \beta \\ 1 & \beta \end{bmatrix}$ is equal to

(a) $\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$

(b) $\begin{bmatrix} -1 & -1 \\ -1 & 2 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$

(d) $\begin{bmatrix} -1 & -1 \\ -1 & -2 \end{bmatrix}$

Ans: (b)

Q) What is the value of the following determinant?

$$\begin{vmatrix} \cos C & \tan A & 0 \\ \sin B & 0 & -\tan A \\ 0 & \sin B & \cos C \end{vmatrix}$$

- (a) -1
- (b) 0
- (c) $2 \tan A \sin B \sin C$
- (d) $-2 \tan A \sin B \sin C$

Q) What is the value of the following determinant?

$$\begin{vmatrix} \cos C & \tan A & 0 \\ \sin B & 0 & -\tan A \\ 0 & \sin B & \cos C \end{vmatrix}$$

- (a) -1
- (b) 0
- (c) $2 \tan A \sin B \sin C$
- (d) $-2 \tan A \sin B \sin C$

Ans: (b)

Q) The value of the determinant $\begin{vmatrix} 1 - \alpha & \alpha - \alpha^2 & \alpha^2 \\ 1 - \beta & \beta - \beta^2 & \beta^2 \\ 1 - \gamma & \gamma - \gamma^2 & \gamma^2 \end{vmatrix}$

is equal to

- (a) $(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma)$ (b) $(\alpha - \beta)(\beta - \gamma)(\gamma - \alpha)$
(c) $(\alpha - \beta)(\beta - \gamma)(\gamma - \alpha)(\alpha + \beta + \gamma)$
(d) 0

Q) The value of the determinant $\begin{vmatrix} 1 - \alpha & \alpha - \alpha^2 & \alpha^2 \\ 1 - \beta & \beta - \beta^2 & \beta^2 \\ 1 - \gamma & \gamma - \gamma^2 & \gamma^2 \end{vmatrix}$

is equal to

- (a) $(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma)$ (b) $(\alpha - \beta)(\beta - \gamma)(\gamma - \alpha)$
(c) $(\alpha - \beta)(\beta - \gamma)(\gamma - \alpha)(\alpha + \beta + \gamma)$
(d) 0

Ans: (b)

Q) What is the value of the determinant

$$\begin{vmatrix} 1! & 2! & 3! \\ 2! & 3! & 4! \\ 3! & 4! & 5! \end{vmatrix} ?$$

(a) 0

(b) 12

(c) 24

(d) 36

Q) What is the value of the determinant

$$\begin{vmatrix} 1! & 2! & 3! \\ 2! & 3! & 4! \\ 3! & 4! & 5! \end{vmatrix} ?$$

(a) 0

(b) 12

(c) 24

(d) 36

Ans: (c)

Q) The inverse of a matrix A is given

$$\text{by } \begin{bmatrix} -2 & 1 \\ 3 & -1 \\ 2 & 2 \end{bmatrix}$$

What is A equal to?

(a) $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & -2 \\ -3 & 4 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 2 \\ 3 & -4 \end{bmatrix}$

(d) $\begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}$

Q) The inverse of a matrix A is given

$$\text{by } \begin{bmatrix} -2 & 1 \\ 3 & -1 \\ 2 & 2 \end{bmatrix}$$

What is A equal to?

(a) $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & -2 \\ -3 & 4 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 2 \\ 3 & -4 \end{bmatrix}$

(d) $\begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}$

Ans: (a)

Q) If $x + a + b + c = 0$, then what is the

value of
$$\begin{vmatrix} x+a & b & c \\ a & x+b & c \\ a & b & x+c \end{vmatrix} ?$$

- (a) 0 (b) $(a + b + c)^2$
(c) $a^2 + b^2 + c^2$ (d) $a + b + c - 2$

Q) If $x + a + b + c = 0$, then what is the

value of
$$\begin{vmatrix} x + a & b & c \\ a & x + b & c \\ a & b & x + c \end{vmatrix} ?$$

- (a) 0 (b) $(a + b + c)^2$
(c) $a^2 + b^2 + c^2$ (d) $a + b + c - 2$

Ans: (a)

Q) What is the value of $\begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}$ if $a^3 + b^3 + c^3 = 0$?

(a) 0

(c) $3 abc$

(b) 1

(d) $-3 abc$

Q) What is the value of $\begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}$ if $a^3 + b^3 + c^3 = 0$?

(a) 0

(c) $3 abc$

(b) 1

(d) $-3 abc$

Ans: (c)

Q) If $1, \omega, \omega^2$ are the cube roots of unity, then

$$\Delta = \begin{vmatrix} 1 & \omega^n & \omega^{2n} \\ \omega^n & \omega^{2n} & 1 \\ \omega^{2n} & 1 & \omega^n \end{vmatrix} \text{ is equal to}$$

- (a) ω^2 (b) 0 (c) 1 (d) ω

Q) If $1, \omega, \omega^2$ are the cube roots of unity, then

$$\Delta = \begin{vmatrix} 1 & \omega^n & \omega^{2n} \\ \omega^n & \omega^{2n} & 1 \\ \omega^{2n} & 1 & \omega^n \end{vmatrix} \text{ is equal to}$$

- (a) ω^2 (b) 0 (c) 1 (d) ω

Ans: (b)

Q) Let A be an $n \times n$ matrix. If $\det(\lambda A) = \lambda^s \det(A)$, what is the value of s ?

(a) 0

(b) 1

(c) -1

(d) n

Q) Let A be an $n \times n$ matrix. If $\det(\lambda A) = \lambda^s \det(A)$, what is the value of s ?

(a) 0

(b) 1

(c) -1

(d) n

Ans: (d)

Q) If a matrix A is such that

$$3A^3 + 2A^2 + 5A + I = 0,$$

Then what is A^{-1} equal to?

- (a) $-(3A^2 + 2A + 5)$ (b) $3A^2 + 2A + 5I$
(c) $3A^2 - 2A - 5I$ (d) $(3A^2 + 2A - 5I)$

Q) If a matrix A is such that

$$3A^3 + 2A^2 + 5A + I = 0,$$

Then what is A^{-1} equal to?

- (a) $-(3A^2 + 2A + 5I)$ (b) $3A^2 + 2A + 5I$
(c) $3A^2 - 2A - 5I$ (d) $(3A^2 + 2A - 5I)$

Ans: (a)

Q) If A is a square matrix, then what is $\text{adj}(A^{-1}) - (\text{adj } A)^{-1}$ equal to?

- (a) $2|A|$
- (b) Null matrix
- (c) Unit matrix
- (d) None of the above

Q) If A is a square matrix, then what is $\text{adj}(A^{-1}) - (\text{adj } A)^{-1}$ equal to?

- (a) $2|A|$
- (b) Null matrix
- (c) Unit matrix
- (d) None of the above

Ans: (b)

Q) Consider the following in respect of the matrix

$$A = \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}:$$

1. $A^2 = -A$

2. $A^3 = 4A$

Which of the above is/are correct?

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Q) Consider the following in respect of the matrix

$$A = \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}:$$

1. $A^2 = -A$

2. $A^3 = 4A$

Which of the above is/are correct?

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Ans: (b)

Q) Let $A = \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$. The only correct

statement about the matrix A is

- (a) $A^2 = I$
- (b) $A = (-1)I$, where I is a unit matrix
- (c) A^{-1} does not exist
- (d) A is a zero matrix

Q) Let $A = \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$. The only correct

statement about the matrix A is

- (a) $A^2 = I$
- (b) $A = (-1)I$, where I is a unit matrix
- (c) A^{-1} does not exist
- (d) A is a zero matrix

Ans: (a)

Q) If $a_1, a_2, a_3, \dots, a_n, \dots$ are in G.P., then the value of the determinant

$$\begin{vmatrix} \log a_n & \log a_{n+1} & \log a_{n+2} \\ \log a_{n+3} & \log a_{n+4} & \log a_{n+5} \\ \log a_{n+6} & \log a_{n+7} & \log a_{n+8} \end{vmatrix}, \text{ is}$$

- (a) -2 (b) 1 (c) 2 (d) 0

Q) If $a_1, a_2, a_3, \dots, a_n, \dots$ are in G.P., then the value of the determinant

$$\begin{vmatrix} \log a_n & \log a_{n+1} & \log a_{n+2} \\ \log a_{n+3} & \log a_{n+4} & \log a_{n+5} \\ \log a_{n+6} & \log a_{n+7} & \log a_{n+8} \end{vmatrix}, \text{ is}$$

- (a) -2 (b) 1 (c) 2 (d) 0

Ans: (d)

Q) The system of equations

$$\alpha x + y + z = \alpha - 1$$

$$x + \alpha y + z = \alpha - 1$$

$$x + y + \alpha z = \alpha - 1$$

has infinite solutions, if α is

(a) -2

(b) either -2 or 1

(c) not -2

(d) 1

Q) If A and B are square matrices of size $n \times n$ such that

$$A^2 - B^2 = (A - B)(A + B),$$
 then which of the following will

be always true?

- (a) $A = B$
- (b) $AB = BA$
- (c) either of A or B is a zero matrix
- (d) either of A or B is identity matrix

Q) If A and B are square matrices of size $n \times n$ such that

$$A^2 - B^2 = (A - B)(A + B),$$
 then which of the following will

be always true?

- (a) $A = B$
- (b) $AB = BA$
- (c) either of A or B is a zero matrix
- (d) either of A or B is identity matrix

Ans: (b)

Q) Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$, $a, b \in N$. Then

- (a) there cannot exist any B such that $AB = BA$
- (b) there exist more than one but finite number of B's such that $AB = BA$
- (c) there exists exactly one B such that $AB = BA$
- (d) there exist infinitely many B's such that $AB = BA$

Q) Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$, $a, b \in N$. Then

- (a) there cannot exist any B such that $AB = BA$
- (b) there exist more than one but finite number of B's such that $AB = BA$
- (c) there exists exactly one B such that $AB = BA$
- (d) there exist infinitely many B's such that $AB = BA$

Ans: (d)

Q) Let $A = \begin{vmatrix} 5 & 5\alpha & \alpha \\ 0 & \alpha & 5\alpha \\ 0 & 0 & 5 \end{vmatrix}$. If $|A^2| = 25$, then $|\alpha|$ equals

(a) $1/5$

(b) 5

(c) 5^2

(d) 1

Q) Let $A = \begin{vmatrix} 5 & 5\alpha & \alpha \\ 0 & \alpha & 5\alpha \\ 0 & 0 & 5 \end{vmatrix}$. If $|A^2| = 25$, then $|\alpha|$ equals

- (a) $1/5$ (b) 5 (c) 5^2 (d) 1

Ans: (a)

Q) Let A be a 2×2 matrix

Statement -1 : $\text{adj}(\text{adj } A) = A$

Statement -2 : $|\text{adj } A| = |A|$

- (a) Statement-1 is true, Statement-2 is true.
Statement-2 is not a correct explanation for Statement-1.
- (b) Statement-1 is true, Statement-2 is false.
- (c) Statement -1 is false, Statement-2 is true.
- (d) Statement-1 is true, Statement -2 is true.
Statement-2 is a correct explanation for Statement-1.

Q) Let A be a 2×2 matrix

Statement -1 : $\text{adj}(\text{adj } A) = A$

Statement -2 : $|\text{adj } A| = |A|$

- (a) Statement-1 is true, Statement-2 is true.
Statement-2 is not a correct explanation for Statement-1.
- (b) Statement-1 is true, Statement-2 is false.
- (c) Statement -1 is false, Statement-2 is true.
- (d) Statement-1 is true, Statement -2 is true.
Statement-2 is a correct explanation for Statement-1.

Ans: (d)

Q) Let $A = \begin{pmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{pmatrix}$. and $10B = \begin{pmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{pmatrix}$. If B is

the inverse of matrix A , then α is

- (a) 5 (b) -1 (c) 2 (d) -2

Q) Let $A = \begin{pmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{pmatrix}$. and $10B = \begin{pmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{pmatrix}$. If B is

the inverse of matrix A , then α is

- (a) 5 (b) -1 (c) 2 (d) -2

Ans: (a)

Q) Let A and B be two symmetric matrices of order 3.

Statement-1: $A(BA)$ and $(AB)A$ are symmetric matrices.

Statement-2: AB is symmetric matrix if matrix multiplication of A with B is commutative.

- (a) Statement-1 is true, Statement-2 is true; Statement-2 is **not a** correct explanation for Statement-1.
- (b) Statement-1 is true, Statement-2 is false.
- (c) Statement-1 is false, Statement-2 is true.
- (d) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

Q) Let A and B be two symmetric matrices of order 3.

Statement-1: $A(BA)$ and $(AB)A$ are symmetric matrices.

Statement-2: AB is symmetric matrix if matrix multiplication of A with B is commutative.

- (a) Statement-1 is true, Statement-2 is true; Statement-2 is **not a** correct explanation for Statement-1.
- (b) Statement-1 is true, Statement-2 is false.
- (c) Statement-1 is false, Statement-2 is true.
- (d) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

Ans: (a)

Q) If $\alpha, \beta \neq 0$, and $f(n) = \alpha^n + \beta^n$ and

$$\begin{vmatrix} 3 & 1+f(1) & 1+f(2) \\ 1+f(1) & 1+f(2) & 1+f(3) \\ 1+f(2) & 1+f(3) & 1+f(4) \end{vmatrix}$$

$= K(1-\alpha)^2(1-\beta)^2(\alpha-\beta)^2$, then K is equal to:

- (a) 1 (b) -1 (c) $\alpha\beta$ (d) $\frac{1}{\alpha\beta}$

Q) If $\alpha, \beta \neq 0$, and $f(n) = \alpha^n + \beta^n$ and

$$\begin{vmatrix} 3 & 1+f(1) & 1+f(2) \\ 1+f(1) & 1+f(2) & 1+f(3) \\ 1+f(2) & 1+f(3) & 1+f(4) \end{vmatrix}$$

$= K(1-\alpha)^2(1-\beta)^2(\alpha-\beta)^2$, then K is equal to:

- (a) 1 (b) -1 (c) $\alpha\beta$ (d) $\frac{1}{\alpha\beta}$

Ans: (a)

Q) If $A = \begin{bmatrix} 5a & -b \\ 3 & 2 \end{bmatrix}$ and $A \text{ adj } A = A A^T$, then $5a + b$ is equal

to :

(a) 4

(b) 13

(c) -1

(d) 5

Q) If $A = \begin{bmatrix} 5a & -b \\ 3 & 2 \end{bmatrix}$ and $A \text{ adj } A = A A^T$, then $5a + b$ is equal

to :

(a) 4

(b) 13

(c) -1

(d) 5

Ans: (d)

Q) If A and B are square matrices of equal degree, then which one is correct among the followings?

(a) $A + B = B + A$

(b) $A + B = A - B$

(c) $A - B = B - A$

(d) $AB = BA$

Q) If A and B are square matrices of equal degree, then which one is correct among the followings?

(a) $A + B = B + A$

(b) $A + B = A - B$

(c) $A - B = B - A$

(d) $AB = BA$

Ans: (a)

Q) If A is a 3×3 non-singular matrix such that $AA' = A'A$ and $B = A^{-1}A'$, then BB' equals:

- (a) B^{-1} (b) $(B^{-1})'$ (c) $I + B$ (d) I

Q) If A is a 3×3 non-singular matrix such that $AA' = A'A$ and $B = A^{-1}A'$, then BB' equals:

- (a) B^{-1} (b) $(B^{-1})'$ (c) $I + B$ (d) I

Ans: (d)

Q) If $A^2 - A + I = 0$, then the inverse of A is

- (a) $A + I$ (b) A (c) $A - I$ (d) $I - A$

Q) If $A^2 - A + I = 0$, then the inverse of A is

- (a) $A + I$ (b) A (c) $A - I$ (d) $I - A$

Ans: (d)

Q) If $a^2 + b^2 + c^2 = -2$ and

$$f(x) = \begin{vmatrix} 1+a^2x & (1+b^2)x & (1+c^2)x \\ (1+a^2)x & 1+b^2x & (1+c^2)x \\ (1+a^2)x & (1+b^2)x & 1+c^2x \end{vmatrix},$$

then $f(x)$ is a polynomial of degree

- (a) 1 (b) 0 (c) 3 (d) 2

Q) If $a^2 + b^2 + c^2 = -2$ and

$$f(x) = \begin{vmatrix} 1+a^2x & (1+b^2)x & (1+c^2)x \\ (1+a^2)x & 1+b^2x & (1+c^2)x \\ (1+a^2)x & (1+b^2)x & 1+c^2x \end{vmatrix},$$

then $f(x)$ is a polynomial of degree

- (a) 1 (b) 0 (c) 3 (d) 2

Ans: (d)

**REVISION
TOPICS :
(19/08/24)**

- **Limits**
- **Continuity**

NDA 2 2024

LIVE

MATHS

REVISION

CLASS 10

SSBCrack
EXAMS



NAVJYOTI SIR

